



COMPILED AND CIRCULATED BY ARPITA CHAKRABORTY, STATE AIDED COLLEGE
TEACHER, DEPARTMENT OF BOTANY, NARAJOLE RAJ COLLEGE.

Topics:

1. Structure of bacterial endospore
2. Formation of bacterial endospore

Bacterial Endospores:

Microorganisms sense and adapt to changes in their environment. When favoured nutrients are exhausted, some bacteria may become motile to seek out nutrients, or they may produce enzymes to exploit alternative resources. One example of an extreme survival strategy employed by certain low G+C Gram-positive bacteria is the formation of endospores. This complex developmental process is often initiated in response to nutrient deprivation. It allows the bacterium to produce a dormant and highly resistant cell to preserve the cell's genetic material in times of extreme stress.

Endospores can survive in environmental stress that would normally kill the bacterium. These stresses include high temperature, high UV irradiation, desiccation, chemical damage and enzymatic destruction. The extraordinary resistance properties of endospores make them of particular importance because they are not readily killed by many antimicrobial treatments. A variety of different microorganisms form "spores" or "cysts", but the endospores of low G+C Gram-positive bacteria are by far the most resistant to harsh conditions.

Structure of bacterial endospore:

The structure of an endospore can be explained in part by its unique cellular structure. The outer proteinaceous coat surrounding the spore provides much of the chemical and enzymatic resistance. Beneath the coat resides a very thick layer of specialized peptidoglycan called the cortex. Proper cortex formation is needed for dehydration of the spore core, which aids in resistance to high temperature.




Fig: Structure of endospore



**COMPILED AND CIRCULATED BY ARPITA CHAKRABORTY, STATE AIDED COLLEGE
TEACHER, DEPARTMENT OF BOTANY, NARAJOLE RAJ COLLEGE.**

A germ cell wall resides under the cortex. This layer of peptidoglycan will become the cell wall of the bacterium after the endospore germinates. The inner membrane, under the germ cell wall, is a major permeability barrier against several potentially damaging chemicals. The centre of the endospore, the core, exists in a very dehydrated state and houses the cell's DNA, ribosomes and large amounts of dipicolinic acid. This endospore-specific chemical can comprise up to 10% of the spore's dry weight and appears to play a role in maintaining spore dormancy. Small acid-soluble proteins (SASPs) are also only found in endospores. These proteins tightly bind and condense the DNA, and are in part responsible for resistance to UV light and DNA-damaging chemicals. Other species-specific structures and chemicals associated with endospores include stalks, toxin crystals, other additional outer glycoprotein layer called the exosporium.

Endospore Development:

The process of forming an endospore is complex. The model organism used to study endospore formation is *Bacillus subtilis*. Endospore development requires several hours to complete. Key morphological changes in the process have been used as markers to define stages of development. As a cell begins the process of forming an endospore, it divides asymmetrically (Stage II). This results in the creation of two compartments, the larger mother cell and the smaller forespore.




Fig: Different stages of Endospore formation

These two cells have different developmental fates. Intercellular communication systems coordinate cell-specific gene expression through these sequential activation of specialized sigma factors in each of the cells. Next (Stage III), the peptidoglycan in the septum is degraded and the forespore is engulfed by the mother cell, forming a cell within a cell. The activities of the mother cell and



**COMPILED AND CIRCULATED BY ARPITA CHAKRABORTY, STATE AIDED COLLEGE
TEACHER, DEPARTMENT OF BOTANY, NARAJOLE RAJ COLLEGE.**

forespore lead to the synthesis of the endospore-specific compounds, formation of the cortex and deposition of the coat (Stages IV+V). This is followed by the final dehydration and maturation of the endospore (Stages VI+VII). Finally, the mother cell is destroyed in a programmed cell death, and the endospore is released into the environment. The endospore will remain dormant until it senses the return of more favourable conditions. [A sigma factor is a small protein that directs RNA polymerase to specific sites on DNA to initiate gene expression.]

Reference-

1. Fundamentals of Microbiology and Immunology, Ajit Kumar Banerjee, Nirmalya Banerjee, New Central Book Agency(P) Ltd, 2006, ISBN: 81-7381-502-X

(All the above mentioned informations are collected from the above reference and will be solely used for teaching and learning purposes).